

*"There's can be no rainbow
without a little rain"*

CSIR NET – Life Science





Unit 1: Biochemistry

12

**Enzyme Kinetics &
Inhibition**



Order Online and Get
Free Delivery Across India





Points to be covered in this Lecture



Catalytic Strategies



Michalis Menten Kinetics



K_m , K_{cat} and V_{max}



LB Plot, HW Plot, EF Plot



Irreversible Inhibitors



Reversible inhibitors





Dixon plot



Catalytic Strategies:

1. General Acid-Base Catalysis

Amino acid residues	General acid form (proton donor)	General Base form (proton acceptor)
Glu, Asp	$\text{R}-\text{COOH}$	$\text{R}-\text{COO}^-$
Lys, Arg	$\text{R}-\overset{\text{H}}{\underset{\text{H}}{\text{N}^+\text{H}}}$	$\text{R}-\ddot{\text{N}}\text{H}_2$
Cys	$\text{R}-\text{SH}$	$\text{R}-\text{S}^-$
His	$\text{R}-\text{C}=\text{CH}-\text{NH}^+$	$\text{R}-\text{C}=\text{CH}-\text{N}:$
Ser	$\text{R}-\text{OH}$	$\text{R}-\text{O}^-$
Tyr	$\text{R}-\text{C}_6\text{H}_4-\text{OH}$	$\text{R}-\text{C}_6\text{H}_4-\text{O}^-$



Transition State

eg Lysozyme
Trypsin
Chymotrypsin


H^+ donor or acceptor

Specific Acid-Base Catalysis

Water



2. Covalent Catalysis



Non-covalent

Substrate
Intermediate

• Trypsin

• Chymotrypsin


• Hydrolytic Enzyme

• Recombinase


Ser-Asp-His



3. Metal Ion Catalysis : Mg^{2+} , Ca^{2+} , Fe^{2+} neutralize -ve charge of substrate so that transition state can be achieved.



4. Electrostatic Catalysis




- $D = \downarrow$
- Electrostatic interaction = \uparrow
- H-bonding = \uparrow
- betⁿ enzyme & substrate





5. Catalysis through Proximity and Orientation Effects

Substrates held close in space
and in correct orientation




Results in more
efficient reaction pathway





ENZYME KINETICS : *Michelis-Menten Kinectic*

• classical enzyme




Where, S is the substrate

E is the enzyme


ES is the enzyme-substrate complex


P is the product

k_1 , k_{-1} , and k_2 are rate constants



Temp^r = 25°
pH = 7
E = 1 nM





$$V_0 = \frac{V_{max} [S]}{K_m + [S]}$$

Where, V_0 = initial reaction velocity

V_{max} = maximal velocity

K_m = Michaelis constant = $\frac{(k_{-1} + k_2)}{k_1}$

$[S]$ = substrate concentration

Dissociation

Association


$$V_{max} \propto E_{Total}$$

$$V_{max} = k_2 E_T$$

$$V_0 \propto [S]$$

K_m is also some constant
substrate concⁿ.

$$V_0 = \frac{1}{2} V_{max}$$




$$V_0 = V_{max} \frac{[S]}{K_m + [S]}$$

$$\frac{V_0}{V_{max}} = \frac{[S]}{K_m + [S]}$$



Assumptions of Michaelis and Menten Equation



$$\frac{V_0}{V_{max}} = \frac{[S]}{K_m + [S]}$$

1. Initial velocity (V_0) = no product, all rxⁿ will proceed forward

2. $[S] \gg [E]$ $[S] = 0.1 \text{ mM to } 1 \text{ mM} \gg [E] = 1 \text{ nM}$

3. ES dissociation is insignificant : $[ES]$ conc^y remains constant

4. Steady-state assumption: Formation of ES = Dissociation of ES

5. Rate limiting step is K2 or

(slowest)

K_{cat}

Turnover number

$$K_1 = K_{-1} + K_2$$

$$K_{-1} \gg K_2$$

$$V_0 = V_{max} \frac{[S]}{K_m + [S]}$$



What is K_m and K_s ?



- Dissociation constant for ES complex

$$K_m = \frac{\text{Dissociation rate constant}}{\text{Association rate constant}} = \frac{k_{-1} + k_2}{k_1}$$

$$K_s = \frac{k_{-1}}{k_1}$$

- Substrate concentration where $V_o = 0.5 V_{max}$

$$\checkmark \frac{V_o}{V_{max}} = \frac{[S]}{K_m + [S]}$$

Suppose $V_o = \frac{1}{2} V_{max}$

$$\frac{1}{2} = \frac{[S]}{K_m + [S]}$$

$$K_m + 1[S] = 2[S]$$


$$K_m = 2[S] - 1[S]$$

$$K_m = [S]$$

$$\begin{array}{l|l} [S] = K_m & V_o = \frac{1}{2} V_{max} \\ [S] < K_m & V_o < \frac{1}{2} V_{max} \\ [S] > K_m & V_o > \frac{1}{2} V_{max} \end{array}$$

- High K_m means low affinity of enzyme for its substrate
- K_m does not vary with the concentration of enzyme or substrate

$$[S] = \uparrow \quad V_o = \uparrow \quad [E] = \uparrow \quad V_{max} = \uparrow$$



$$E_1 = K_{M2} = 10 \text{ mM}$$

$$E_2 = K_{M2} = 20 \text{ mM}$$

• E_1 has higher affinity for its substrate



Apply your Mind

$$V_0 = V_{max} \frac{(S)}{K_m + (S)}$$

$$V_0 = 100 \frac{\mu\text{mol}}{\text{min}} \times \frac{10 \text{ mM}}{10 \text{ mM} + 10 \text{ mM}}$$

$$V_0 = 100 \times \frac{10}{20} = 50 \frac{\mu\text{mol}}{\text{min}}$$

The enzyme alkaline phosphatase was tested for its catalytic activity using the substrate para-nitrophenylphosphate. The K_m obtained was 10 mM and V_{max} was 100 $\mu\text{mol}/\text{min}$. Which one of the following options represents the initial velocity of the reaction at a substrate concentration of 10 mM?

- (1) 50 $\mu\text{mol}/\text{min}$
- (2) 100 $\mu\text{mol}/\text{min}$
- (3) 500 $\mu\text{mol}/\text{min}$
- (4) 20 $\mu\text{mol}/\text{min}$

$$\begin{aligned} &\checkmark [S] = K_m & V_0 &= 0.5 V_{max} \\ &[S] > K_m & V_0 &> 0.5 V_{max} \\ &[S] < K_m & V_0 &< 0.5 V_{max} \\ &10 \text{ mM} = 10 \text{ mM} & V_0 &= 0.5 \times 100 \frac{\mu\text{mol}}{\text{min}} \\ & & &= 50 \end{aligned}$$



Apply your Mind

$$V_0 = V_{max} \frac{(S)}{1K_m + (S)}$$

What is the fold difference between V_0 at $[S] = \frac{1}{2} K_m$ and V_0 at $[S] = 2K_m$ where V_0 is the initial velocity of an enzyme catalyzed reaction, $[S]$ is substrate concentration and K_m is the Michaelis constant?

(1) 2 Fold

(2) 4 Fold

(3) 6 Fold

(4) 8 Fold

Rx 1

$V_0 = ?$

$[S] = 0.5 K_m$

$$V_0 = V_{max} \frac{0.5 K_m}{1 K_m + 0.5 K_m}$$

$$= V_m \frac{0.5 K_m}{1.5 K_m}$$

$$= V_m \times \frac{1}{3}$$

$$V_0 = 0.33 V_{max}$$

Rx 2

$V_0 = ?$

$[S] = 2 K_m$

$$V_0 = V_{max} \frac{2 K_m}{1 K_m + 2 K_m}$$

$$V_0 = V_{max} \frac{2 K_m}{3 K_m}$$

$$V_0 = \frac{2}{3} V_{max}$$

$$V_0 = 0.66 V_{max}$$



What is K_{cat} or turnover number?

$$V_{max} = K_2 E_T$$

$$K_{cat} = V_{max} / E_T$$

$$V_{max} = K_{cat} E_T$$


collective speed

Speed of single enzyme

Total Enzyme

K_{cat} is more : enzyme is fast
 K_{cat} is less : enzyme is slow

K_{cat} is constant, don't vary with substrate concⁿ or enzyme concⁿ.



$$V_{max} = M / \text{sec}$$

$$E_T = M$$

$$V_0 = \frac{V_{max} (S)}{K_m + (S)}$$



Fast and slow enzymes

i) Enzyme	ii) Turnover number
Catalase ✓	4.0×10^7
Carbonic anhydrase ✓	1.0×10^6
Acetylcholine esterase ✓	2.5×10^4
Lactate dehydrogenase	1.0×10^3
Urease ✓	1.0×10^4
Chymotrypsin ✓	100.0
Rubisco Slow	→ 3.0
Lysozyme Slow	→ 0.5

Per second

K_{cat} or k_2 or

$$\text{Turnover (} K_{cat} \text{)} = \frac{V_{max}}{E_T}$$

$$= \frac{M/sec}{M}$$

$$K_{cat} = 1/sec$$

$$K_{cat} = sec^{-1}$$

First order rate constant


*For Ribozymes and restriction endonuclease, it is expressed

per minute

very slow



Specificity constant or catalytic efficiency




K_{cat}/K_M is rate constant that refers to properties and reactions of free enzyme and free substrate

Enzyme	Substrate	$K_M (M)$	$k_{cat} (s^{-1})$
✓ Acetylcholinesterase	Acetylcholine	9.5×10^{-5}	1.4×10^4
✓ Carbonic anhydrase	CO_2	1.2×10^{-2}	1.0×10^6
✓ Superoxide dismutase	Superoxide ion ($O_2^{\cdot -}$)	3.6×10^{-4}	1.0×10^6
✓ Catalase	H_2O_2	2.5×10^{-2}	1.0×10^7

K_{cat}/K_M

$$1.4 \times 10^4 / 9.5 \times 10^{-5} \approx \dots \times 10^8$$

$$10^6 / 1.2 \times 10^{-2} \approx \dots \times 10^8$$




K_{cat}/K_M of 2×10^8 for enzyme indicates that the value is close to diffusion controlled rate of encounter



Line-weaver-Burk equation and Plot

$$V_0 = \frac{V_{max} (S)}{K_m + (S)}$$



Rectangular Hyperbolic curve

Difficult to obtain V_{max} & K_m precisely by MM plot
convert it into straight line $y = mx + c$

Double Reciprocal of MM eqⁿ to obtain straight line


$$\frac{1}{V_0} = \frac{K_m + [S]}{V_{max} [S]}$$

$$\frac{1}{V_0} = \frac{K_m}{V_{max}} \cdot \frac{1}{S} + \frac{[S]}{V_{max}} \cdot \frac{1}{[S]}$$

$$\frac{1}{V_0} = \frac{K_m}{V_{max}} \cdot \frac{1}{[S]} + \frac{1}{V_{max}}$$

$$y = mx + c$$

Slope = $\frac{K_m}{V_{max}}$



Independent Variable



The Hanes-Woolf plot

LB BURK equation multiplied by $[S]$

$$\frac{[S]}{v_0} = \frac{1}{V_{max}} \cdot [S] + \frac{K_M}{V_{max}}$$

$$Y = m \cdot x + c$$


$$\text{Slope} = \frac{1}{V_{max}}$$

$$\text{Reciprocal of slope} = V_{max}$$

$$\text{x-axis intercept} = K_M$$

$$1/K_M = 1/4$$

$$K_M = 4$$





Eadie-Hofstee Plot (Imp)

The Eadie-Hofstee's equation is derived from Michaelis-Menten equation as shown below,

$$V_0 = \frac{V_{max} [S]}{K_M + [S]}$$

$$V_0 (K_M + [S]) = V_{max} [S]$$

$$V_0 K_M + V_0 [S] = V_{max} [S]$$


Divide it by $[S]$

$$\frac{V_0 K_M}{[S]} + \frac{V_0 [S]}{[S]} = \frac{V_{max} [S]}{[S]}$$

$$\frac{V_0}{[S]} \cdot K_M + V_0 = V_{max}$$

$$V_0 = -K_M \cdot \frac{V_0}{[S]} + V_{max}$$

$$Y = -m \cdot x + C$$





Expression of enzyme activity

✓ 1 UNIT:

one μmole of substrate transformed or product formed per Minute under defined conditions ($\mu\text{mole}/\text{min}$)



$$1 \text{ unit} = 1 \frac{\mu\text{mole}}{\text{min}}$$

✓ 1 Katal:

$$1 \text{ Katal} = 1 \frac{\text{mole}}{\text{sec}}$$

$$1 \text{ Katal} = 6 \times 10^7 \text{ units}$$

one mole of substrate transformed or product formed per Second under defined conditions





INHIBITION OF ENZYME ACTIVITY

$$\epsilon_T = \downarrow$$

$$V_{max} = \downarrow$$

Irreversible Inhibitor:

Binds covalently (tightly) to enzyme
Enzyme activity is permanently lost.

I. Group-specific covalent modifying agents:**Serine:**

- Diisopropylfluorophosphate (DIFP)
- Phenylmethyl sulphonylfluoride (PMSF).

Cysteine:

Iodoacetate
Iodoacetamide
N-ethylmaleimide
 Ag^+ , Hg^{2+} , Pb^{2+}

eg **Trypsin**
Chymotrypsin

Protease cleaves after lysine or arginine

eg Caspase
Glyceraldehyde 3-Phosphate
dehydrogenase



2. Affinity labels: binds to substrate binding site covalently

Tosyl Phe~~ky~~l~~al~~anyl Chloromethyl Ketone (TPCK):

Binds covalently to histidine in substrate binding site of chymotrypsin.

protease

→ cleaves after
aromatic aa (Phe, Tyr, Trp)

3. Transition state analogs: resembles transition state of substrate
& binds covalently to substrate binding site



Pyrrole 2- carboxylate which resemble with transition state of proline and inhibits enzyme proline racemase.



4. Suicide inhibitors or k_{cat} inhibitors

Eg. Allopurinol \rightarrow Xanthine oxidase

Inhibitor directly don't bind to enzyme but if inhibitor are modified by enzyme than modified form binds covalently to enzyme.






B. Reversible Inhibitors:

Binds non-covalently to enzyme



1. Competitive inhibition : Substrate & Inhibitor both compete for same substrate binding site.





ES or EI binary complex formation
 • Inhibition can also be reversed by increasing substrate conc^r.

Effect on V_{max} : unchanged

Effect on K_m : Increase, Affinity decrease



Effect on MM and Lineweaver-Burke plot:






Examples of competitive inhibitors:

- **Statin group** drugs inhibits **HMG CoA reductase** (cholesterol biosynthesis)
- **Malonic acid** inhibits **succinate dehydrogenase**
- **Sulpha drugs** (sulfonamide) inhibits **dihydropteroate synthase (DHPS)** $\xrightarrow{\text{folic acid biosynthesis}}$
- **Methotrexate** and **aminopterin** inhibits **dihydrofolate reductase**
 $\text{H}_2\text{-folate} \xrightarrow{\quad} \text{H}_4\text{-folate (THF)}$
- **ddi** and **AZT** inhibit **reverse transcriptase**
- **Ethanol** inhibit **alcohol dehydrogenase**

\hookrightarrow Treatment of methanol poisoning



2. Non-competitive inhibition




Effect on V_{max} : Decrease

Effect on K_m : No change, affinity of enzyme for substrate enzyme remain same



Effect on MM and Lineweaver-Burke plot:





Examples of non-competitive inhibitors:

Cyanide inhibits cytochrome oxidases

complex IV of electron transport chain






3. Mixed inhibition

Binding site of (S) & (I) are different.

Binding of (I) decrease affinity for (S) = $K_m \uparrow$




- Binary (EI) & Ternary (ESI) complex are formed

Effect on V_{max} : Decrease

Effect on K_m : Increase




Effect on MM and Lineweaver-Burke plot:





Example: Phosphoenolpyruvate carboxykinase Inhibited by Genistein




mixed inhibitor

$K_m = \uparrow$
 $v_{max} = \downarrow$



4. Uncompetitive inhibitors





Effect on V_{max} : **Decrease**
 Effect on K_m : **Decrease**

- Binding site for substrate & inhibitor are different.
- Inhibitor cannot bind to free enzyme
No EI complex
- Presence of inhibitor favors binding of substrate (affinity = \uparrow)
 $K_m = \text{Decrease}$
- Once ES complex forms, inhibitor binding site get exposed & inhibitor also binds to ES complex to form ESI ternary complex



Effect on MM and Lineweaver-Burke plot:





Examples of uncompetitive inhibitors:

L- Phenylalanine inhibits *Alkaline phosphatase*




Inhibition	Lineweaver Burk Equation
A. Competitive ✓	$1. \frac{1}{v_o} = \left(\frac{\alpha K_M}{V_{\max}} \right) \frac{1}{[S]} + \frac{1}{V_{\max}}$ <p><i>Handwritten: $y = mx + c$</i></p>
<u>B. Un-competitive</u>	$2. \frac{1}{v_o} = \left(\frac{K_M}{V_{\max}} \right) \frac{1}{[S]} + \frac{\alpha'}{V_{\max}}$ <p><i>Handwritten: Slope same</i></p>
C. Mixed	$3. \frac{1}{v_o} = \left(\frac{\alpha K_M}{V_{\max}} \right) \frac{1}{[S]} + \frac{\alpha'}{V_{\max}}$



Dixon Plot

Important

Plotting between the reciprocal of the reaction velocity ($1/v$) against various concentrations of an inhibitor ($[I]$) at a fixed substrate (s) concentration.



$$[s] = 0.1 \text{ mM}$$

$$[s] = 0.2 \text{ mM}$$

$$\text{x-axis} = [I]$$

← Independent

$$\text{y-axis} = 1/v_0$$

← Dependent

x-axis intercept

$$K_i$$



Not important

Competitive Inhibition:


Lines intersect on the x-axis (inhibitor concentration axis).

Non-competitive Inhibition:

Lines intersect on the y-axis (reciprocal velocity axis).

Uncompetitive Inhibition:

Lines are parallel (no intersection).





Apply Your Mind

Following statements are made about uncompetitive inhibition of an enzyme:

- A. Inhibitor binds to free enzyme but not to enzyme- substrate complex. ✗
- ✓ B. Addition of uncompetitive inhibitor lowers the apparent V_{max} and apparent K_m of the reaction.
- ✗ C. Apparent K_m of the enzyme is increased but V_{max} remains same
- ✓ D. Inhibitor binds to free enzyme- substrate complex but not to free enzyme

Which one of the following option represents the correct combination of the statements ?

- (1) B and C
- (2) ✗ A and C
- (3) B and D ✓
- (4) ✗ A and D

③



Thank you

Learn From India's Best Educators
India's **No.1** EdTech Company for Graduates
& Post Graduates Examination.

➡ Download **IFAS** App Now



ANDROID



IOS



WINDOWS



Harshada
Sharma

Earned:
₹25,500

"As is widely recognized, IFAS stands as India's leading institute for CSIR NET examination preparation. My Personal philosophy has always revolved around helping others and I've had the pleasure of introducing the IFAS Online Course to 12 of my classmates and juniors, which in turn allowed me to earn 25,500 rupees. You too can join in and refer your friends, allowing you to earn a 5% rewards based on the fees each of your referred friends Pay"



Step – 01 Invite Friends

Share your referral code or Link with your friends.

Step – 02 Earn Cashback

On Every Purchase made using your referral code, you earn 5% instant cashback in your bank account.

Step – 03 Refer More, Earn More

Keep referring as many friends as you can and keep **Earning** cash as much as you can



Friendship is Sharing, Caring, Giving & Helping Each Other...!